

## Diagnostic accuracy of organ electrodermal diagnostics

### A pilot study

J Szopinski, D Pantanowitz, G G Jaros

**Objective.** To estimate the diagnostic accuracy as well as the scope of utilisation of a new bio-electronic method of organ diagnostics.

**Design.** Double-blind comparative study of the diagnostic results obtained by means of organ electrodermal diagnostics (OED) and clinical diagnoses, as a criterion standard.

**Setting.** Department of Surgery, Helen Joseph Hospital, Johannesburg.

**Patients.** 70 pre-selected inpatients of mean age 36 (SD = 7) years with suspected pathology of one (or more) of the following organs: oesophagus, stomach, duodenum, biliary tract, pancreas, colon, kidneys and urinary tract. In total, 276 of the abovementioned internal organs were selected for statistical consideration.

**Main outcome measures.** The difference between the so-called basic electrical impedance of the skin and the impedance value established for a particular organ projection area (the skin zone corresponding to a particular internal organ).

**Results.** In total 250 true OED results were obtained from the 276 subjects considered: detection rate 90.6% (95% CI 87.1 - 94.1%). Established OED sensitivity was 91.8% (95% CI 88.6 - 95.0%) and OED specificity equalled 89.9% (95% CI 86.4 - 93.4%). The predictive value for positive OED results was 83.3% (95% CI 78.9 - 87.7%) and for negative OED results 95.2% (95% CI 92.0 - 98.4%). The OED results were affected neither by the type nor the aetiology of disease, i.e. OED estimates the actual extent of the pathological process within particular organs but does not explain the cause of pathology directly. No side-effects of the OED examinations were observed.

**Conclusions.** So-called organ projection areas do exist on the skin surface. The electrical impedance of the

projection areas corresponding to diseased organs is increased, relative to that of healthy organ-related skin zones. The difference in impedance is proportional to the intensity of the pathological process. OED, which utilises these electrical phenomena of the skin, may detect diseased organs and estimate the extent of pathological process activity within these organs.

*S Afr Med J* 1998; 88: 146-150.

Various specific relationships between the skin and internal organs are known. Pain sensitivity (e.g. as assessed by means of Head's dermatomes), skin temperature, colour as well as electrical parameters may be changed by internal organ pathology. Correlations between skin electrical resistance and psychological status (psychogalvanic reaction), endocrinological function and autonomic innervation of particular dermatomes are also well known. The electrical current perception threshold demonstrates changes in many diseases.

Many authors have investigated the specific influences of particular organ pathology on the electrical parameters of the corresponding skin areas.<sup>1-11</sup> Diagnostic methods based upon measurements of electrical potential, resistance and impedance of these zones have been proposed; however, their diagnostic accuracy has not been proven and reproducibility has not been consistent. Some of these methods utilise specific bio-electrical properties of acupuncture points.<sup>1,3-11</sup> These include: 'EAV' by Voll<sup>4,5,7,10,11</sup> with modification 'Vegatest',<sup>4</sup> 'Ryodoraku' after Nakatani<sup>1,3,7,10</sup> with modification 'CITO',<sup>7,10</sup> many types of auricular electropunctural diagnostics<sup>6,7,10</sup> and some methods based upon corporeal acupuncture point impedance measurements.<sup>7,8,10</sup>

Different measurement techniques and different electrical parameters of the measurement currents have been used in the abovementioned methods. The results obtained often depend on the patient's muscular tension, emotional condition, skin humidity, procedure duration, environmental temperature and the pressure of the measuring electrode. Therefore these methods did not find widespread application in contemporary medicine and the authors' ideas did not create a unified and systematised scientific basis for the utilisation of bio-electrical skin properties for organ diagnostics.

The first author's own investigations of the skin electrical characteristics<sup>9,12-16</sup> showed that after the so-called electrical breakthrough effect was obtained in skin areas related to diseased organs, an asymmetry of electrical resistance characteristics was observed. This asymmetry is based upon the increased skin resistance for a positively polarised measuring electrode compared with the resistance for negative polarisation of the same electrode (semi-conductivity phenomenon). Also, the impedance of skin zones corresponding to diseased organs is increased, compared with that of the healthy organs' projection areas.

The abovementioned findings create a basis for a new non-invasive diagnostic method — organ electrodermal diagnostics (OED).<sup>9,17-23</sup> The location of electrical resistance asymmetry and increased impedance areas on the skin surface, although often remote, may indicate which organs

Departments of Family Medicine and Surgery, Coronation and Helen Joseph hospitals and University of the Witwatersrand, Johannesburg

J Szopinski, DM

D Pantanowitz, BSc, MB BCH, FRCS, FCS

Department of Biomedical Engineering, University of Cape Town

G G Jaros, MSc (Eng), PhD (Present address: Department of Anaesthetics, University of Sydney, Australia)

are involved in the pathological process. The degree of asymmetry or the difference in impedance may indicate the activity of disease.

This research study is aimed at estimating the diagnostic accuracy as well as the scope of utilisation of a new OED system, 'Diagnotronics'.

## Patients and methods

A comparative study of clinical diagnoses and OED results was performed by means of a double-blind trial on a group of 70 inpatients at Helen Joseph Hospital's surgical department. The group consisted of 37 men and 33 women of mean age 36 (SD = 7) years. During the post-intake ward rounds, the surgical consultants had pre-selected newly admitted patients with suspected pathology of one (or more) of the following organs: oesophagus, stomach, duodenum, biliary tract, pancreas, colon, kidneys and urinary tract. These organs are relatively easy to access clinically, and their pathologies represent a variety of aetiological and pathogenetic factors, e.g. infections, inflammation, neoplasms, and immunological and metabolic disorders.

In each case, the OED examination of all the abovementioned organs was undertaken before the final clinical diagnosis was established. The OED investigator had no access to the patient's documentation whatsoever and OED examinations were always witnessed independently to avoid the suspicion of any communication between investigator and patient. The documented OED results were placed in a special container immediately after the examination was completed; this was only opened after the routine clinical investigation had been concluded.

Clinical investigations of the chosen internal organs comprised:

1. Oesophagus — history and physical examination, chest radiograph, barium swallow, oesophagoscopy with biopsy for confirmation/exclusion of oesophagitis or a neoplastic process. Operative findings were included if the patient had undergone surgery.
2. Stomach and duodenum — history and physical examination, barium meal, gastroduodenoscopy with biopsy for confirmation/exclusion of mucosal inflammation or a neoplastic process. Operative findings were included if the patient had undergone surgery.
3. Biliary tract — history and physical examination, acute phase indicators, liver function tests, hepatitis markers, urine for bilirubin and urobilinogen assessment, ultrasound examination, cholecystogram/cholangiogram (if indicated), hepatic immuno-diacetic acid (HIDA) cholescintigraphy (if indicated). Operative findings were included, if the patient had undergone surgery.
4. Pancreas — history and physical examination, serum and urine amylase, blood glucose, faecal fats, ultrasound examination, abdominal radiograph, computed tomography (CT) scan, endoscopic retrograde cholangiopancreatography (ERCP). Operative findings were included, if the patient had undergone surgery.
5. Colon — history and physical examination, full blood count, barium enema, sigmoidoscopy and/or colonoscopy. Liver function test, liver ultrasound examination, CT scan (if indicated). Operative findings were included, if the patient

had undergone surgery.

6. Kidneys and urinary tract — history and physical examination, urine for microscopy culture and susceptibility, urea and electrolytes, creatinine clearance, acute phase indicators, ultrasound examination, intravenous pyelogram. CT scan, cystoscopy and renal biopsy were performed if indicated. Operative findings were included, if the patient had undergone surgery.

OED examinations were performed by means of the prototype OED device, Diagnotronics (Fig. 1). This electronic apparatus makes organ diagnoses based on an automatic impedance evaluation of the skin areas corresponding to particular internal organs. To avoid the problem of individual basic skin impedance, which is different for each person, Diagnotronics first estimates the impedance value for a calibration electrode of 2 cm diameter; this is used as a point of reference and then compared with the diagnostic results obtained with a measurement electrode of 1 mm diameter.<sup>20</sup>



Fig. 1. OED by means of Diagnotronics.

The examination entails placement of the reference electrode on any area of the patient's skin, e.g. on a hand, and the placement of the measurement electrode, after the abovementioned calibration, on the skin area corresponding to the particular organ (organ projection areas of ear auricles were used in this study) (Fig. 2). Using a liquid crystal display, the OED device specifies the actual condition of the organ related to the investigated skin area as 'Healthy 0°', 'Disease I°' (within normal limits), 'Disease II°' (subacute pathology), 'Disease III°' (acute pathology). A special display graded according to percentage of the disease intensity makes it possible to specify accurately the activity of organ pathology. By means of a TV screen the device demonstrates the locations of skin areas corresponding to the examined organs; it also projects diagnoses after the examination is completed.

A special OED information sheet was given to each patient prior to examination. Ethical approval was obtained from the Ethics Committee of the University of the Witwatersrand.

The comparison of clinical diagnoses and OED results was undertaken by an independent arbiter who was not




Fig. 2. Location of chosen organ projection areas on ear auricle: 1 - oesophagus; 2 - stomach; 3 - duodenum (left auricle); 4 - gall bladder (left auricle); 5 - pancreas (right auricle); 6 - kidney; 7 - colon; 8 - urinary bladder.

involved in the diagnostic procedures. OED sensitivity was defined as the proportion of correctly classified positives, i.e. 'true positives' among the total of diseased persons.<sup>24,25</sup> The sensitivity rate was estimated with the formula:

$$s = \frac{a}{(a + c)} \times 100$$

s = sensitivity rate; a = true positive; c = false negative.

OED specificity was defined as the proportion of true negatives among the total of persons free of disease.<sup>24,25</sup> The specificity rate was estimated with the formula:

$$sp = \frac{d}{(b + d)} \times 100$$

sp = specificity rate; d = true negative; b = false positive.

The positive predictive value of the OED result was defined as the probability of having the disease among the group of persons classified as positive by OED:<sup>24,25</sup>

$$PV (pos) = \frac{a}{(a + b)} \times 100$$

PV = predictive value rate; a = true positive; b = false positive.

The negative predictive value of the OED result was defined as the probability of not having the disease among the group of persons classified as negative by OED:<sup>24,25</sup>

$$PV (neg) = \frac{d}{(c + d)} \times 100$$

PV = predictive value rate; d = true negative; c = false negative.

The  $\chi^2$ -test was used to calculate statistical significance.  $P < 0.05$  was accepted as the statistically significant difference. Only organs with proven clinical conditions (healthy/diseased) were considered for final statistical comparison.

## Results

In total, 250 true OED results were obtained from the 276 subjects considered: detection rate 90.6% (95% CI 87.1 - 94.1%) (Table I). Established OED sensitivity was 91.8% (95% CI 88.6 - 95.0%) and OED specificity equalled 89.9% (95% CI 86.4 - 93.4%). The predictive value for positive OED results was 83.3% (95% CI 78.9 - 87.7%) and for negative OED results 95.2% (95% CI 92.0 - 98.4%).

The patient subgroups representing one of the chosen internal organs were generally too small for statistical conclusions (pilot study) but some tendencies were noted: healthy organs usually display the OED result 'Healthy' or 'Disease I', while subacute pathology displays 'Disease II' and acute pathology 'Disease III'. The OED results were affected neither by the type nor the aetiology of disease, i.e. OED estimated the actual extent of pathological process activity within particular organs but did not explain the direct cause of pathology.

The OED results were not influenced by a patient's muscular tension, emotional state, skin humidity, environmental temperature, or by procedure duration. The pressure of the measuring electrode had a limited influence (up to 5%) on the OED results and did not affect final diagnoses. No side-effects of the OED examinations were observed.

## Discussion

OED probably belongs to that area of medical diagnostics which utilises the nervous system's information function.<sup>26</sup> The OED diagnostic results obtained in this study confirmed the existence of so-called organ projection areas on the skin surface. These specific areas have been used especially in reflexive physiotherapy, e.g. acupuncture, pressopuncture, analgesic electrostimulation, laser therapy and reflexive thermotherapy. However, there was no direct evidence of the real functional connection between the skin surface and particular internal organs. OED has made possible the precise individual localisation of optimal skin areas for reflexive physiotherapy, which might increase its therapeutic efficacy.

In comparison with other known diagnostic methods which utilise the connection between bio-electrical skin properties and the state of internal organs,<sup>1-11</sup> OED results are fully reproducible and reliable. They are independent of procedure duration and the patient's muscular tension,

**Table 1. Comparison of clinical diagnoses and OED results obtained by means of the Diagnetronics device**

Organ	Clinical diagnosis	No. of subjects	True OED results						False OED results					
			Negative			Positive			Negative			Positive		
			0°	I°	Together	II°	III°	Together	0°	I°	Together	II°	III°	Together
Oesophagus	Healthy	26	13	12	25							1		1
	Oesophagitis	8				4	3	7	1	1				
	Cancer	6					6	6						
Stomach	Healthy	18	4	11	15							2	1	3
	Gastritis	15				8	4	12	1	2	3			
	Ulcers	3				1	2	3						
	Cancer	4				1	3	4						
Duodenum	Healthy	26	8	14	22							3	1	4
	Duodenitis	5				4	1	5						
	Ulcers	4					3	3	1	1				
Gall bladder	Healthy	27	8	17	25							2		2
	Gallstone	15				5	8	13	2	2				
	Cholecystitis													
	Acute	2					2	2						
	Chronic	3				2	1	3						
Pancreas	Healthy	7	2	3	5							1	1	2
	Pancreatitis													
	Acute	3					3	3						
Colon	Healthy	4	4		4									
	Colitis	4				3	1	4						
	Cancer	3				2	1	3						
Kidneys	Healthy	41	20	18	38							3		3
	Pyelonephritis	2				2		2						
	Nephrolithiasis	7				1	6	7						
Urinary bladder	Healthy	29	17	9	26							3		3
	Cystitis	1					1	1						
	Cancer	4				2	2	4						
Total		276			160*			90*			8			18

\* Statistically significant difference between the total sum of true and false results:  $P < 0.0001$ .

Detection rate = 90.6% (87.1 - 94.1%); Sensitivity rate = 91.8% (88.6 - 95.0%); Specificity rate = 89.9% (86.4 - 93.4%); Predictive value rate (pos.) = 83.3% (78.9 - 87.7%); Predictive value rate (neg.) = 95.2% (92.0 - 98.4%).

emotional state, skin humidity, and environmental temperature, and show only a limited correlation with the measuring electrode's pressure on the skin.

This pilot study indicates the high diagnostic accuracy of OED as well as its usefulness in clinical practice. A full clinical evaluation of this new bio-electronic system of non-invasive medical diagnostics should now be undertaken with an adequate number of patients.

## Conclusions

1. So-called organ projection areas do exist on the skin surface. The electrical impedance of the projection areas corresponding to diseased organs is increased, relative to healthy organ-related skin zones. The difference in impedance is proportional to the intensity of the pathological process.

2. OED, which utilises the abovementioned electrical

phenomenon of the skin, seems to be a reliable bio-electronic method of non-invasive medical diagnostics, with high rates of sensitivity and specificity. OED may detect diseased organs and estimate the extent of pathological process activity within these organs.

3. The OED results are affected neither by the type nor the aetiology of disease, i.e. OED cannot directly explain the cause of pathology.

The authors express their gratitude to the Bioenergy Association (chaired by Dr B Brom) for the financial support of this research project.

## REFERENCES

- Garnuszewski Z. *Renaissance of Acupuncture* (Polish). Warsaw: 'Sport i turystyka', 1988.
- Glaser-Tuerk M. Spezielle lokale Tests zum Nachweis aktiver Herde im Kieferbereich. *Oester Z Stom* 1968; 65(6): 222-226.
- Hyodo MD. *Ryodoraku Treatment and Objective Approach to Acupuncture*. Osaka: 1975.
- Kenyon JN. Diagnostic techniques using bioenergetic recording methods: The science of bioenergetic regulatory medicine. *Am J Acupuncture* 1986; 14(1): 5-13.
- Köhnlechner M. *Handbuch der Naturheilkunde*. Munich: Kindler Verlag, 1975.

6. Nogier P. *Traite d'Auriculotherapie*. Moulins-les Metz: Maisonneuve, 1988.
7. Portnov F. *Electropuncture Reflexotherapy* (Russian). Riga: 'Zinatne' Publishing House, 1982.
8. Rosenblatt S. The electrodermal characteristics of acupuncture points. *Am J Acupuncture* 1982; 10(2): 131-137.
9. Szopinski J. The use of bioelectrical properties of skin for organ diagnostics (Polish). *Wiad Lek* 1989; 42: 697-702.
10. Tabayeva D. *Manual of Acupuncture Reflexotherapy* (Russian). Moscow: Meditsina, 1980.
11. Voll R. *Topographische Lage der Messpunkte der Elektroakupunktur* Uelzen: Dr Blume & Co; ML Publishers, 1988.
12. Clazek M, Szopinski J, Skrzypulec V. Investigations of morphological structure of acupuncture points and meridians. *J Tradit Chin Med* 1985; 5(4): 289-292.
13. Szopinski J, Sierak T, Gabryel A. Die Erforschung der Bioenergetischen Eigenschaften der Akupunkturmeridiane. *Dtsch Z Akup* 1988; 2: 31-35.
14. Szopinski J, Sierak T, Kanlewski M, Niezbecki A. Der Einfluss ausgewählter Krankheiten auf die Bioelektrischen Eigenschaften der Akupunkturpunkte. *Dtsch Z Akup* 1988; 3: 51-55.
15. Szopinski J, Sierak T, Niezbecki A, Kanlewski M. Influence of selected internal diseases on electrical parameters of acupuncture points (Polish). In: Sapinski W, ed. *Diary of 1st National Conference on Acupuncture*, 23 - 24 September 1982. Warsaw: Kallaz: WSZ, 1982; 54-69.
16. Szopinski J, Sierak T, Szopinska H, Clazek M. Bioelektrische und Bioenergetische Eigenschaften und morphologische Strukturen der Akupunkturpunkte und Akupunkturmeridiane. In: Bischko J, ed. *Handbuch der Akupunktur und Auriculotherapie*. Part 25.2.0. Heidelberg: Haug Verlag, 1985; 105-124.
17. Sierak T, Szopinski J. Universal device for Electrodermal Organ Diagnostics and electrotherapy (Polish). *Probl Tech Med* 1987; 18(4): 255-262.
18. Sierak T, Szopinski J. Universelles elektronisches Gerät für die automatische Elektropunkturdiagnostik und Elektroreflextherapie. *Dtsch Z Akup* 1988; 6: 112-115.
19. Szopinski J. Application of the Organ Electrodermal Diagnostics own method in chosen internal diseases (Polish). PhD thesis. Silesian Medical Academy, Katowice; 1985.
20. Szopinski J. An impedance measuring apparatus. Provisional patent No. 95/5790. RSA, 1995.
21. Szopinski J, Macura T, Sierak T, et al. Experimental determination of skin projection areas of liver and their use in monitoring of viral hepatitis course. *Clin Phys Physiol Meas* (in press).
22. Szopinski R, Sierak B, Sierak R, Pastor A. The device for a measurement of the electrical resistance asymmetry of acupuncture points. Patent No. 146183. Poland, 1989.
23. Szopinski J, Sierak T, Sztklinarz J, et al. Experimental determination of skin projection areas of stomach and duodenum and their use in monitoring of ulcer disease course. *Clin Phys Physiol Meas* (in press).
24. Fletcher RH, Fletcher SW, Wagner EH. *Clinical Epidemiology. The Essentials*. Baltimore: Williams & Wilkins, 1991.
25. Toman K. Sensitivity, specificity and predictive value of diagnostic tests. *Bull Int Union Tuberc* 1981; 56: 18-27.
26. Szopinski J, Sierak T, Wilk A. Neurophysiological foundation of the Organ Electrodermal Diagnostics, acupuncture and other methods of the reflexive physiotherapy. *Am J Acupuncture* (in press).

Accepted 3 Oct 1997.

# Human bites of the face

## A review of 22 cases

Midion M Chidzonga

**Objective.** To compare experience with human bites of the face in a unit in Harare, Zimbabwe, with other similar studies in respect of reasons for the assaults, age and sex of the patients, timespan between injury and treatment, surgical management and incidence of infection.

**Design.** Prospective study.

**Setting.** Department of Oral and Maxillofacial Surgery Outpatients Clinic, Harare Central Hospital, Harare, Zimbabwe.

**Patients.** Twenty-two consecutive patients with human bites of the face.

**Main outcome measures.** The black female is the predominant victim and assailant, with the lower lip most commonly involved.

**Results.** Of the patients 81.8% were female. The mean age was 32.5 years. Interpersonal violence was the commonest cause of the injury, with the lower lip involved in 90.9% of cases. Early presentation reduces the risk of infection.

**Conclusion.** The results indicate that the black female predominates both as victim and assailant. Early presentation with thorough surgical debridement under antibiotic cover produces satisfactory results.

*S Afr Med J* 1998; 88: 150-152.

Human bites usually occur during quarrels, but they can also occur during the course of sexual activity.<sup>1,2</sup> The latter form of bite is uncommon in our practice, possibly because kissing as a sign of affection is not as common among black Africans as it is in Caucasians.<sup>1,3,4</sup>

Human bites of the face are serious injuries, mainly because of the potential for severe wound infection, disability or gross disfigurement.<sup>3,5,6</sup> The main objectives of management are therefore prevention of infection and functional closure of the wound with the best possible cosmetic result.<sup>5</sup>

The present study in our unit in Harare, Zimbabwe, comprises 22 human bites of the face (defined by Tomasetti *et al.*<sup>6</sup> as 'one inflicted on a person by another person'). The main aim of the study was to compare the results with other similar studies in respect of incidence of infection, surgical management, age and sex of the patients, cause of the injuries and timespan between injury and treatment.